



Sensitive periods in development and risk for psychiatric disorders and related endpoints: a systematic review of child maltreatment findings

Jonathan D Schaefer, Theresa W Cheng, Erin C Dunn

Lancet Psychiatry 2022;
9: 978–91

Department of Psychology,
University of Minnesota,
Minneapolis, MN, USA
(J D Schaefer PhD); Psychiatric
and Neurodevelopmental
Genetics Unit, Center for
Genomic Medicine,
Massachusetts General
Hospital, Boston, MA, USA
(T W Cheng PhD, E C Dunn ScD);
Department of Psychiatry,
Harvard Medical School,
Boston, MA, USA (T W Cheng,
E C Dunn); Center on the
Developing Child, Harvard
University, Cambridge, MA,
USA (E C Dunn)

Correspondence to:
Dr Jonathan D Schaefer,
Department of Psychology,
University of Minnesota,
Minneapolis, MN 55455, USA
schae567@umn.edu

Variation in the mental health of people who have experienced childhood maltreatment is substantial. One hypothesis is that this variation is attributable, in part, to the timing of maltreatment—specifically, whether maltreatment occurs during sensitive periods in development when the brain is maximally sensitive to particular types of environmental input. To determine whether there is scientific consensus around when periods of peak sensitivity occur, we did a systematic review of human observational studies. Although 89 (75%) of the 118 unique cross-sectional or longitudinal cohort studies we identified reported timing effects, no consistent sensitive periods were identified for any of the most studied outcomes. Thus, observational research on childhood maltreatment has yet to converge on a single period (or set of periods) of increased vulnerability. We identified study characteristics that might contribute to these between-study differences and used observations from our Review to suggest a comprehensive set of recommendations for future research.

Introduction

Psychiatric disorders affect an estimated 16–26% of the global population each year and 70–80% of the population at some point during their lifetime.^{1–3} A major driver of psychiatric disorders is childhood maltreatment, a form of early adversity that encompasses the absence of expected inputs and responses from parents or caregivers (eg, physical or emotional neglect) and the presence of harmful, threatening inputs and responses (eg, physical, sexual, or emotional abuse).^{4,5} Childhood maltreatment is common, with worldwide prevalence estimates ranging from 13% (sexual abuse) to 36% (emotional abuse). People with a history of maltreatment have, on average, at least twice the risk of developing a psychiatric disorder,⁶ an effect that appears to be causal.^{7,8} Nevertheless, it is clear that not all maltreated children are equally affected. One important determinant of the effects of maltreatment might be its developmental timing—specifically, whether it occurs during sensitive periods in development, meaning life stages when particular environmental inputs might be more influential. Identifying when these periods exist could have major benefits, including advancing understanding of the mechanisms linking maltreatment to later psychopathology and informing the development of targeted strategies to facilitate more timely intervention.

Neuroplasticity, which refers to the brain's dynamic capacity to undergo structural and functional maturation and change in response to experience,⁹ is a powerful determinant of the extent to which different life experiences can become biologically embedded. So-called critical and sensitive periods are states of elevated plasticity when the effect of experience on neurodevelopment is especially strong and lasting.^{5–7} Critical periods have long been viewed from a deterministic perspective: after the critical period ends, the effects of experience are permanent and irreversible; the window of plasticity closes and remains shut. But windows of sensitive-period

plasticity do not fully close; rather, some degree of plasticity remains, although considerably more effort is required to produce change after a sensitive period ends than when that window was open. Therefore, we favour the term sensitive period in this Review, because of the potential for symptomatic improvement, remission, or recovery in nearly all common forms of psychopathology.

The neurobiological mechanisms that create sensitive periods are complex and encompass mechanisms that inhibit, initiate, and later slow, plasticity.¹⁰ These mechanisms involve processes that govern the initial wiring of the brain (eg, synapse formation and overproduction) and those that shape consolidation of neural networks (eg, synapse elimination and pruning).⁹ The associated psychological literature also commonly groups these mechanisms into those that are time limited and activated by universal experiences across the human species (sometimes labelled as experience expectant), and other ongoing learning mechanisms thought to be unique to the individual and their environment (ie, experience dependent).¹¹ The experience-expectant category is more commonly implicated in the formation of sensitive periods; however, the distinction between the two types of mechanisms is not necessarily precise and this framework does not capture all types of plasticity.¹²

Scientific interest in sensitive periods started nearly a century ago, when the concept was first articulated by Charles Stockard, an embryologist. Stockard observed that birth defects in fish offspring occurred only when mothers were exposed to extreme temperatures and toxic chemicals just as a specific embryonic organ was developing, labelling these time periods as moments of supremacy or critical moments.¹³ In 1935, ethologist Konrad Lorenz was the first to apply this concept to animal behaviour, noting that young ducklings, goslings, and chicks would not imprint in the typical way for their species if their encounter with a moving object was delayed.¹⁴ In the second half of the 20th century, animal

researchers also began generating evidence to support the existence of critical socialisation periods when specific social inputs were necessary for typical social and emotional functioning.¹⁵ Harry Harlow's 1965 study¹⁶ showed rhesus monkeys kept in social isolation between ages 3 months and 12 months had clear, enduring social and emotional deficits (whereas isolation at later ages appeared not to have such an effect), and in 1970, Nobel laureates David Hubel and Torsten Wiesel began establishing critical periods in the development of sensory systems.¹⁷ These early animal socialisation studies, in particular, informed human research on attachment and approach learning, and the hypothesis that early caregiving relationships have a crucial role in determining later emotional and behavioural health.¹⁸

Since the 1950s, studies mentioning sensitive or critical periods have steadily increased. Articles additionally mentioning maltreatment are fewer but increased by 40 times in less than two decades (appendix 1 p 4). Published reviews investigating timing effects for maternal deprivation or institutionalisation suggest a sensitive period between ages 6 months and 24 months.¹⁹ However, periods of peak vulnerability to maltreatment remain unknown.^{20,21}

In this systematic review, we investigated whether evidence has converged to identify sensitive periods when the developing brain is most susceptible to the effects of childhood maltreatment. Specifically, we investigated the questions: (1) what proportion of studies testing for timing effects of maltreatment find such effects, and how consistent are the periods of peak vulnerability across studies? (2) Some researchers have postulated that sensitive periods are a property of individual biological systems (eg, neural circuits) rather than complex phenotypes, such as mental health disorders;²² are findings more consistent in studies at more "biological" levels of analysis? (3) Do findings become more consistent when studies are stratified by different study characteristics (eg, sample size, statistical approach, or assessment methods)? (4) Research has repeatedly suggested a high degree of multifinality in psychopathology, wherein the same risk factor increases risk for multiple types of disorder simultaneously;²³ does this principle extend to studies of the timing effects of maltreatment (ie, do multiple disorder types show similar timing effects)? (5) Relatedly, the principle of equifinality suggests that the same mental health problems can arise through multiple different developmental histories and risk factors;²³ does this principle extend to studies of timing effects (ie, do different types of maltreatment show similar timing effects)? (6) Are there sex or gender differences in the periods identified across studies?

Methods

Search strategy and selection criteria

Using PRISMA guidelines,²⁴ we did a systematic review of human observational studies that tested for

maltreatment timing effects by searching MEDLINE, PsycINFO, and the Cumulative Index to Nursing and Allied Health. Because some researchers have speculated that sensitive periods might be more readily identified for more biological outcomes, we sought to identify studies examining whether timing of maltreatment was related to psychopathology (eg, psychiatric disorder diagnosis or high scores on symptom scales) as well as studies connecting timing of maltreatment to related constructs within other levels of analysis, including neuroimaging (ie, brain structure or functioning), epigenetics (ie, DNA methylation), psychophysiology (eg, cortisol reactivity or cortisol awakening response), and behaviour (eg, personality or cognitive test performance). In addition to covering multiple levels of analysis, we selected search terms to capture a range of Research Domain Criteria constructs, including negative valence systems, positive valence systems, cognitive systems,

See Online for appendix 1





Figure 1: Study profile

CINAHL=Cumulative Index to Nursing and Allied Health.



social processes, arousal and regulatory systems, and sensorimotor systems.²⁵

The inclusion criteria used in this Review were that studies reported results from human participants; the research was published in English; participants were assessed for one or more relevant outcomes; participants were assessed for exposure to maltreatment alone or alongside other measures of adversity; maltreatment exposure was assessed for at least two non-overlapping time periods that each preceded the outcome assessment (studies assessing maltreatment exposure across a continuous range of ages were considered to meet this criterion); and the article reported results for at least one test of sensitive period effects, whether that was an implicit (ie, non-statistical) or explicit comparison of associations between maltreatment occurring during two or more different age periods and a given outcome. Studies were labelled as being of “maltreatment” if they assessed any of these exposures: physical abuse, sexual abuse, emotional or verbal abuse, physical neglect, or emotional neglect. Although we required studies to assess maltreatment occurring at ages before age 18 years, we did not exclude studies for also assessing maltreatment after this point or based solely on participant age (because many were retrospective studies of adults). These criteria and definitions were established before doing our systematic search; however, the search procedure was not preregistered and a public protocol was not prepared.

Our search proceeded in four steps. First, we searched each database from inception to Aug 9, 2020, using strings related to child abuse, maternal deprivation, neglect, sexual, physical, or emotional abuse, maltreatment, adverse childhood experiences, and victimization, as well as strings related to age, timing, and sensitive or critical periods, to psychopathology or mental disorders, to other Research Domain Criteria such as negative or positive valence, and to neuroimaging; full strings are shown in appendix 1 (pp 5–9). We exported

articles to the systematic review application, Rayyan.²⁶ Second, two authors (JDS and ECD) independently screened titles and abstracts of articles retrieved from the search before reviewing the text of potentially eligible studies. Third, two authors (JDS and ECD) screened the text of retrieved articles to identify those meeting all inclusion criteria. Fourth, two authors (JDS and ECD) independently extracted study characteristics and our primary outcome (the sensitive periods identified). Discrepancies at each step were resolved by discussion. We arranged studies into subgroups based on key characteristics, including the outcome of interest, maltreatment type assessed, and other study design features, and then we searched the references of all included studies.

We evaluated all studies for risk of bias with two independent raters (two of three raters for each study: JDS, TWC, and Samantha Stoll) using an adapted Quality Assessment Tool for observational cohort and cross-sectional Studies developed by the National Heart, Lung, and Blood Institute.²⁷ Discrepancies were resolved by the third rater. Finally, we used permutation testing to statistically evaluate sensitive periods for the most studied outcomes in each level of analysis (ie, depression, language or cognitive ability, and hippocampal volume; appendix 1 pp 2–3).

Results

Our search identified 10830 studies, of which 114 were eligible to be included and a further four studies were identified through citation searches, such that 118 independent reports met our inclusion criteria (figure 1). Of these studies, 71 (60%) tested for associations of maltreatment timing with psychopathology, 8 (7%) with neural indices, 5 (4%) with DNA methylation, 25 (21%) with other Research Domain Criteria units of analysis, and 9 (8%) examined outcomes at multiple levels (appendix 2 p 1).

Despite a shared focus on maltreatment, the 118 studies differed substantially. First, sample sizes were highly variable, ranging from 15 to 978 647 participants. 100 (85%) studies included more than 100 participants, with 37 (31%) reporting more than 1000 participants (figure 2A).

Second, maltreatment exposure was measured with reports of individual maltreatment types, maltreatment clusters, and maltreatment experiences co-occurring with other childhood adversities (figure 2B). Only 28 (24%) of the 118 studies employed prospective assessments; the remainder used cross-sectional or retrospective designs. 61 (52%) studies used measures that aggregated across different maltreatment types or combined maltreatment with other kinds of adversity and 50 (42%) examined only specific maltreatment types (most commonly sexual abuse). The remaining 7 (6%) studies used both aggregated and specific measures.

Figure 2: Overview of the literature on sensitive periods for child maltreatment

Full details on the 118 studies are in appendix 2. (A) Scatterplot of increases in sample size across sensitive period studies. (B) Frequency of child maltreatment exposures examined across sensitive period studies. Studies could be counted more than once if they examined multiple maltreatment types. Maltreatment composite studies combined multiple types of maltreatment into a single exposure measure, and adversity composite studies combined maltreatment and other non-maltreatment adversities into a single exposure measure. (C) Frequency of specific outcomes examined across sensitive period studies. Studies could be counted more than once if they examined multiple outcomes. (D) Use of exposure timing, including dividing age into two to six categories or examining year of age (continuous). (E) Frequency of analytical strategies used in sensitive period studies. Indirect comparison studies described rather than tested differences in the magnitude of the association between exposure and outcome as a function of exposure timing; direct comparison studies statistically evaluated whether the effect of exposure at one point differs from that at another; and competing hypotheses studies directly evaluated alternative theoretical explanations, including accumulation, chronicity, or recency. GMV=grey matter volume.


See Online for appendix 2

Third, several dozen outcomes were studied; psychopathology was the most common. Within our three major outcome categories (psychopathology, imaging, and other Research Domain Criteria units of analysis), the most common outcomes were depression, hippocampal volume, and language or cognitive ability (figure 2C).

Fourth, researchers defined maltreatment timing differently. Definitions of timing included the age at first or worst exposure; the presence or absence of maltreatment during different periods of development (eg, age 0–11 years vs age 12–18 years); scores showing the extent of exposure during different periods of development (eg, frequency, severity, total number of maltreatment events, or number of types of maltreatment); and grouping by recency of maltreatment, often combined with the onset of maltreatment (eg, early maltreatment vs early-and-recent or recent-only maltreatment). Timing was most often treated as a categorical measure, with development

divided into periods roughly corresponding to early childhood, middle childhood, and adolescence. Considerably fewer studies used dichotomous or continuous measures of age (figure 2D).

Fifth, studies varied in the analytical rigour with which they identified sensitive periods. The least rigorous type of study, which we labelled as indirect comparison studies (n=41 [35%]), described rather than statistically tested differences in the magnitude of the association between maltreatment and outcome as a function of exposure timing. Direct comparison studies (n=55 [47%]) provided a stronger test of the sensitive period hypothesis by statistically evaluating whether the effect of exposure at one point in development differed from the effect of exposure at another. The most rigorous type, competing hypotheses studies (n=20 [17%]), directly compared the effects of maltreatment happening during two or more



(Figure 3 continues on next page)

time periods and evaluated alternative explanations for observed timing effects, including maltreatment accumulation (number of exposure types or times exposed), chronicity (duration), or recency (only timing of the most recent exposure). Two (2%) of the 118 studies did not fit into any of these categories (figure 2E).

Our assessment of study quality showed that the risk of biased results from this literature was high overall (appendix 2 p 2). Across studies, the mean number of study quality criteria met was 5.5 (SD 1.9) of 13. The most common item that did not pass the risk of bias assessment was whether the paper presented a justification for selecting or recruiting the number of participants included or discussed statistical power. Only 21 (18%) studies included this information, of which 15 (71%) indicated they were likely to be underpowered for at least some of the analyses done.

Across all 118 studies, 89 (75%) reported at least one sensitive period. Although sensitive periods appeared more consistent when exposure age was dichotomised, no consistent sensitive periods were identified for each of the most commonly studied outcomes in each major level of analysis (depression, hippocampal volume, and language or cognitive ability; figure 3). This observation was supported with permutation testing, which indicated insufficient evidence to reject the null hypothesis that childhood maltreatment was not associated with the outcome at each age (appendix 1 p 11). No clear timing patterns were observed among the next most studied outcomes within each level, although the number of studies within these domains was too small for permutation testing (appendix 1 pp 12–13).

Although some experts have suggested that sensitive periods might be more readily found for more “biological” outcomes,²² we observed only weak evidence supporting this hypothesis. For example, whereas neuroimaging and epigenetic studies were more likely to report sensitive period effects than studies of psychopathology (8 [100%] of the 8 neuroimaging studies were positive vs 4 [80%] of 5 epigenetics studies vs 54 [76%] of 71 psychopathology studies), these differences did not reach statistical significance, which was probably due to the small number of studies in some groups ($\chi^2=2.45$, $df=2$, $p=0.293$).

There was also no clear relationship between likelihood of reporting positive findings and analytical rigour. The proportion of studies reporting at least one sensitive period was statistically comparable across competing hypotheses studies and indirect comparison studies (18 [90%] of 20 competing hypotheses studies and 35 [85%] of 41 indirect comparison studies; $\chi^2=0.25$, $df=1$, $p=0.615$). However, direct comparison studies reported significantly fewer positive findings (35 [64%] of 55) than both competing hypotheses studies ($\chi^2=4.918$, $df=1$, $p=0.027$) and indirect comparison studies ($\chi^2=5.616$, $df=1$, $p=0.018$).

Because of the range of factors that could contribute to the mixed findings we observed, stratifying papers by




Figure 3: Summary of maltreatment-related sensitive period findings for depressive symptoms or depression diagnosis (A), hippocampal volume (B), and cognitive or language ability (C), by timing as dichotomous, categorical, or continuous age

Each line shows the results from one study,^{83–127} with the length and position of the line showing the ages when maltreatment was assessed. Thicker, colour-coded line segments indicate the specific sensitive periods identified by the study. Solid dots show the end of the maltreatment exposure assessment window. Open dots indicate studies in which maltreatment exposure was assessed beyond age 18 years. Coloured gradients show sensitive periods from studies using age as continuous and reporting that outcomes were associated with an earlier or later sensitive period without denoting specific ages.

outcome alone might be insufficient to investigate between-study heterogeneity. Therefore, we tried organising papers on depression (the most common study outcome) in other ways—including by sample size, analytical rigour, maltreatment assessment approach, study design, and use of diagnosis versus symptom outcomes—to see whether more consistent findings would emerge (figure 4). They did not. Similarly, no clear patterns emerged when grouping studies according to timing of the outcome assessment (ie, whether depression was measured in adolescence, young adulthood, or in heterogeneous age groups; appendix 1 p 14).

We tested whether multiple types of mental health problems might share similar sensitive periods by reviewing studies assessing internalising and externalising disorders. Although these outcomes overlap, they are more causally and symptomatically distinct than disorders within the same diagnostic group (eg, anxiety and depression). Although we found substantial between-study variation in the sensitive periods observed for both internalising and externalising disorders, individual studies tended to find similar periods of peak vulnerability for both outcomes (appendix 1 p 15).


To investigate whether different maltreatment types share similar sensitive periods, we identified 20 studies that tested sensitive period effects for at least two maltreatment types. Of these studies, 15 (75%) reported at least one positive sensitive period finding, with all but two (both of which used a dichotomous measure of age) also finding that sensitive periods differed at least somewhat across maltreatment type (figure 5). However, there was little consistency across these studies in the sensitive periods identified for specific types of maltreatment (eg, physical vs sexual abuse) or broader maltreatment categories (eg, abuse vs neglect).

Sex differences in the timing of sensitive periods could also be expected, because of known sex differences in maltreatment prevalence and timing,^{4,83} brain development,⁸⁴ psychopathology risk,^{85,86} and sex-specific critical periods for hormonal and corticolimbic system functioning in rodents.^{87,88} The few studies examining sex differences were almost equally split on whether sex differences in the effects of timing were found (appendix 1 pp 16–17).

Discussion

Although three-quarters of the 118 studies included in this systematic review reported timing effects, we found little converging evidence for maltreatment-related sensitive periods. Sensitive periods were not more consistent in studies with similar characteristics, including outcome type, operationalisation of maltreatment timing, or study rigour.

Possible explanations for why we observed such low convergence include that the timing and magnitude of sensitive periods has been shown to vary as a function of both genes and environmental input (this variability probably complicates efforts to identify universal periods of enhanced neuroplasticity),^{86,89} or that sensitive periods in childhood are less numerous and their effects more nuanced than those occurring prenatally or perinatally. Periods of neuroplasticity that occur later in development might therefore be considerably more difficult to identify in observational work.



(Figure 4 continues on next page)

However, neither of these explanations would explain why our maltreatment results contrast those found in studies examining maternal deprivation or institutionalisation, which suggest larger effects on future psychopathology during the first 6–24 months of life.²⁰ This discrepancy might partly be because maternal deprivation and institutionalisation both represent omissions of

expected care, whereas the category of maltreatment is more heterogeneous, encompassing both omission of expected care (ie, neglect) and unexpected harm (ie, abuse). However, the literature on deprivation and institutionalisation is characterised by considerable complexity, with some reports indicating a second sensitive period in pre-adolescence,⁹⁰ and both




Figure 4: Maltreatment-related sensitive period studies for depressive symptoms or depression diagnosis, by sample size (A), analytical strategy (B), maltreatment assessment approach (C), study design (D), and diagnosis versus symptom outcomes (E)

Each line shows the results from one study,⁸³⁻¹²⁷ with the length and position of the line showing the ages when maltreatment was assessed. Thicker line segments indicate the specific sensitive periods identified by the study. Solid dots show the end of the maltreatment exposure assessment window. Open dots indicate studies in which maltreatment exposure was assessed beyond age 18 years. Coloured gradients show sensitive periods from studies with age as continuous and reporting that outcomes were associated with an earlier or later sensitive period.




Figure 5: Sensitive periods identified by studies assessing two or more different types of maltreatment and psychopathology outcomes, by timing as dichotomous, categorical, or continuous age
 Results of each study¹²⁸⁻¹³⁷ are shown as a group of coloured lines, with the length and position of the lines showing the ages at which maltreatment was assessed and colour showing maltreatment type. Thicker segments indicate the specific sensitive periods identified for each outcome, whereas thinner segments indicate no sensitive periods were identified for that age range. Solid dots show the end of the exposure assessment window. Gradients show sensitive periods from studies using age as continuous and reporting that worse outcomes were associated with an earlier or later sensitive period. Results are shown only from studies with at least one positive sensitive period finding.

sex-specific and sleeper (or latent) effects.^{8,91,92} Even studies using randomised assignment to study the effects of institutionalisation have been unable to disentangle the effects of exposure duration from developmental timing.¹⁹ These sources of complexity also extend to studies of maltreatment, perhaps further contributing to inconsistent findings across studies.

Another possible contributor, cited in nearly all systematic reviews, is methodological heterogeneity. We attempted to understand this heterogeneity by grouping studies with similar characteristics. However, these efforts were hindered in many instances by the small number of studies in some subsets. The possibility of heterogeneity that was unaccounted for even within these subsets makes it difficult to discern if conflicting findings indicate a true absence of sensitive periods or simply methodological and statistical confounds. At the very least, our findings suggest that if maltreatment timing effects exist, they are likely to be sensitive to aspects of study design. In the panel, we outline some of the most probable contributors to these discrepancies and steps to mitigate them in future research.

We found that 75% of studies assessing children for multiple types of maltreatment reported distinguishable sensitive periods associated with each type. This observation aligns with theories organising exposures along distinct continuums of adversity that differentially affect neural systems and behaviour (eg, the Dimensional Model of Adversity and Psychopathology).^{93,94} Thus, our findings suggest sensitive periods might be clarified by examining specific forms of maltreatment or dimensions of adversity rather than broad or aggregated composites.

Our results additionally suggest a need for increased precision in assessments of maltreatment timing. Dichotomous operationalisations of age seemed to contribute to a false sense of consistent findings, which disappeared when more precise measures of maltreatment timing were used. Studies with more precise measures of timing using instruments such as the Maltreatment and Abuse Chronology of Exposure scale⁸³ might help to identify relatively brief (eg, <3 years) sensitive periods that would otherwise be missed.⁹⁵⁻⁹⁷

Studies using objective outcome measures (eg, brain morphometry or stress physiology) were not statistically more likely to identify sensitive periods than studies using subjective outcomes (eg, psychiatric symptoms). However, the small number of high-quality studies using objective outcomes combined with heterogeneity in the outcomes studied limited our ability to make strong inferences about whether shifting focus to more transdiagnostic, biological measures would better show maltreatment timing effects.

Regarding subjective outcomes, we observed that individual studies assessing both internalising and externalising symptoms generally identified overlapping sensitive periods for these constructs. This finding might reflect within-study correlations of these

outcomes, and also suggests that sensitive periods relevant to maltreatment could manifest as increased vulnerability to the development of general psychopathology, rather than specific disorders or diagnostic families. This explanation aligns with decades of research indicating that the effects of maltreatment are non-specific, increasing risk for multiple types of psychiatric disorders simultaneously.^{7,98,99} Consequently, our understanding of maltreatment timing effects might benefit from first examining outcomes at the level of general psychopathology, unless there are compelling theoretical reasons to examine disorder-specific measures.

Risk of bias was high in the studies we reviewed, with many not meeting even half of the criteria in our quality assessment tool. One salient limitation was that few studies reported being adequately statistically powered. This finding suggests it would be premature to interpret the null or mixed findings summarised in this systematic review as strong evidence that there are no sensitive period effects. Future studies on sensitive periods should address this limitation by estimating statistical power and preregistering analytical plans on open science platforms to reduce concerns about multiple testing. Studies should also try to assess and communicate how much the reported frequency of a particular exposure (eg, sexual abuse) varies across development, because this variation could affect power to detect sensitive periods at specific ages (eg, early in development when infantile amnesia limits maltreatment recall).

Only 24% of the studies had a prospective assessment strategy. Although sensitive periods for depression from prospective studies were not more reliable than from retrospective studies, known challenges with retrospective measures suggest their predominance is a limitation of the field. Compared with prospective measures, retrospective measures typically overestimate maltreatment effects on subjective outcomes and underestimate maltreatment effects on objective outcomes.^{100,101} Retrospective and prospective measures of maltreatment occurrence also show low levels of agreement.¹⁰² To our knowledge, agreement between retrospective and prospective measures of maltreatment timing has not been systematically studied, but it might be even lower due to cognitive biases, including telescoping effects (the tendency to perceive recent events as more remote and more distant events as more recent).¹⁰³ We therefore encourage investigators examining maltreatment timing effects to use prospective assessment strategies whenever possible.

In addition to more rigorous measurement of maltreatment, we encourage innovation in the development of new and more objective measures of early-life stressors, as these could reduce measurement error and increase statistical power to detect timing effects. For example, combining self-report and parent-report measures of maltreatment with glucocorticoid levels

Panel: Key challenges and recommendations for future research on maltreatment timing effects

- Timing effects probably differ for different types of maltreatment, which peak in frequency and severity at different ages and work through partly separable neural mechanisms. Researchers should therefore stratify maltreatment analyses by maltreatment type or broad dimensions of adversity (eg, deprivation vs threat) when possible
- Over-reliance on crude measures of age of exposure (eg, dichotomised into early vs late) might obscure brief sensitive periods and contribute to a false sense of consistency that disappears when more precise measures of age are used. Researchers should therefore avoid dichotomous operationalisations of age in favour of multicategorical or continuous options
- The effects of maltreatment on psychopathology are likely to be general (ie, increasing risk of multiple types of mental health problems) rather than specific, and yet most sensitive period studies examine disorder-specific outcomes. Unless compelling reasons exist to examine disorder-specific measures, researchers should consider first examining outcomes at the level of general psychopathology or broad psychiatric spectra (eg, internalising symptoms)
- Risk of biased findings appears high because many studies are of poor quality. Researchers can take steps to improve quality by adopting the changes described, and ensuring adequate statistical power and preregistering analyses
- Over-reliance on retrospective measures of maltreatment, which show poor agreement with prospective measures, typically overestimate maltreatment effects on subjective outcomes, and might generate biased estimates of maltreatment timing. Researchers should therefore use prospective assessment strategies with frequent assessments whenever possible, and consider incorporating objective measures of exposure (eg, biomarkers)
- Observational studies of sensitive periods face statistical challenges, including multicollinearity (because maltreatment exposure tends to be highly correlated across adjacent ages), multiple testing (when numerous age ranges or regression models are compared), and many ways to operationalise relevant constructs. Researchers should therefore consider using advanced analytical approaches including the Structured Life Course Modeling Approach, random forest regression, or specification curve analysis
- Ruling out alternative explanations is crucial; what appears to be developmental differences in plasticity could instead reflect myriad other age-related differences (eg, rates of exposure or child-coping resources). Although no study will be able to account for all potential alternative explanations, researchers could increase confidence in their findings by testing as many plausible competing hypotheses as their data allow

sampled repeatedly across development might help researchers to account partly for some of the known issues within the literature on maltreatment (eg, recall failure and cognitive biases).¹⁰⁴ Children's primary teeth have also recently been proposed as a promising new tool, because of their unique ability to record the timing of early-life stressors at a daily and weekly resolution.¹⁰⁵ Teeth might well be a new and non-invasive biospecimen that records life experiences objectively and can guide causally informative research on how and why psychiatric disorders emerge after experiences of adversity.¹⁰⁶

When assessing sensitive periods in observational studies, two key challenges exist: multicollinearity and multiple testing. Multicollinearity arises because maltreatment exposure can be highly correlated across adjacent

ages. Multiple testing problems arise when several age ranges or regression models are compared. To mitigate these issues, new analytical strategies have been developed to test sensitive period effects with accumulation-of-risk and other theories, including the structured life course modelling approach^{107,108} and machine-learning approaches (eg, random forest regression).¹⁰⁹ In addition, specification curve analysis can address methodological heterogeneity issues, because this technique examines how effect sizes vary across hundreds or even thousands of models that operationalise key parameters (eg, maltreatment type or timing) in different ways.¹¹⁰

Eliminating alternative explanations for apparent sensitive period effects is crucial. What appears as developmental differences in plasticity could instead reflect age-related differences in rates, severity, or accumulation of maltreatment exposure, or other factors that vary by age, such as child coping resources and cognitive capacity (eg, younger children might internalise maltreatment because they do not have the cognitive or socioemotional sophistication to attribute it to parental characteristics). Ruling out all alternative explanations is only possible in an experimental context. Nevertheless, investigators with observational data could increase confidence in their findings by testing as many plausible competing hypotheses as possible.

Although theories on the importance of developmental timing exist in multiple fields, including life-course epidemiology,¹¹¹ developmental neuroscience,²² and developmental psychopathology,¹¹² this systematic review shows that these theories are not yet consistently supported by evidence from human observational studies on childhood maltreatment. We hope this Review inspires more rigorous research to investigate these timing effects, which, if known, could inform the optimal timing of intervention efforts aimed at improving mental health at the individual and population levels.

Conclusions

Researchers and clinicians should be cautious before deeming a particular phase of development a sensitive period for the effects of childhood maltreatment on psychopathology and associated constructs, because existing research does not align on any single period of increased vulnerability. This observation should not be interpreted as conclusive evidence that maltreatment-related sensitive periods do not exist, because it could also be due to methodological heterogeneity, study designs characterised by high risk of bias, and low analytical rigour. Studies that directly address these limitations are needed before information regarding maltreatment timing can be productively incorporated into causal models and clinical decision making.

Contributors

JDS and ECD did the literature search; JDS made the tables and figures; JDS, TWC, and ECD all contributed to data analysis and interpretation, and the writing and editing of the manuscript.

Declaration of interests

JDS received funding from the National Institute on Drug Abuse (award number T32DA050560). ECD received funding from the National Institute of Mental Health of the National Institutes of Health (award number R01MH113930). TWC declares no competing interests.

Acknowledgments

We thank Susan T Landry for her assistance in copyediting the manuscript and Dr Andrew Simpkin for consultation on statistical methods, as well as Janine Cerutti, Emma Glickman, Olivia Sterns, Samantha Stoll, and Amy Riegelman for their assistance in literature searches and data extraction for this Review. The sources providing funding for each of the authors played no role in the study design; collection, analysis, and interpretation of data; writing of the report; or decision to submit the paper for publication. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

References

- 1 Kessler RC, Chiu WT, Demler O, Merikangas KR, Walters EE. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry* 2005; **62**: 617–27.
- 2 Schaefer JD, Caspi A, Belsky DW, et al. Enduring mental health: prevalence and prediction. *J Abnorm Psychol* 2017; **126**: 212–24.
- 3 Rehm J, Shield KD. Global burden of disease and the impact of mental and addictive disorders. *Curr Psychiatry Rep* 2019; **21**: 10.
- 4 Stoltenborgh M, Bakermans-Kranenburg MJ, Alink LRA, van IJzendoorn MH. The prevalence of child maltreatment across the globe: review of a series of meta-analyses. *Child Abuse Rev* 2015; **24**: 37–50.
- 5 Barnett D, Manly J, Cicchetti D. Defining child maltreatment: the interface between policy and research. *Child Abuse Child Dev Soc Policy*. 1993; **8**: 7–73.
- 6 Gilbert R, Widom CS, Browne K, Fergusson D, Webb E, Janson S. Burden and consequences of child maltreatment in high-income countries. *Lancet* 2009; **373**: 68–81.
- 7 Schaefer JD, Moffitt TE, Arseneault L, et al. Adolescent victimization and early-adult psychopathology: approaching causal inference using a longitudinal twin study to rule out noncausal explanations. *Clin Psychol Sci* 2018; **6**: 352–71.
- 8 Wade M, Fox NA, Zeanah CH, Nelson CA. Effect of foster care intervention on trajectories of general and specific psychopathology among children with histories of institutional rearing: a randomized clinical trial. *JAMA Psychiatry* 2018; **75**: 1137–45.
- 9 Ismail FY, Fatemi A, Johnston MV. Cerebral plasticity: windows of opportunity in the developing brain. *Eur J Paediatr Neurol* 2017; **21**: 23–48.
- 10 Reh RK, Dias BG, Nelson CA 3rd, et al. Critical period regulation across multiple timescales. *Proc Natl Acad Sci USA* 2020; **117**: 23242–51.
- 11 Nelson CA 3rd, Gabard-Durnam LJ. Early adversity and critical periods: neurodevelopmental consequences of violating the expectable environment. *Trends Neurosci* 2020; **43**: 133–43.
- 12 Frankenhuis WE, Walasek N. Modeling the evolution of sensitive periods. *Dev Cogn Neurosci* 2020; **41**: 100715.
- 13 Stockard CR. Developmental rate and structural expression: an experimental study of twins, 'double monsters' and single deformities, and the interaction among embryonic organs during their origin and development. *Am J Anat* 1921; **28**: 115–277.
- 14 Lorenz K. Der Kumpan in der Umwelt des Vogels. *J Ornithol* 1935; **83**: 289–413.
- 15 Scott JP, Marston MV. Critical periods affecting the development of normal and mal-adjustive social behavior of puppies. *J Genet Psychol* 1950; **77**: 25–60.
- 16 Harlow HF, Dodsworth RO, Harlow MK. Total social isolation in monkeys. *Proc Natl Acad Sci USA* 1965; **54**: 90–97.
- 17 Hubel DH, Wiesel TN. The period of susceptibility to the physiological effects of unilateral eye closure in kittens. *J Physiol* 1970; **206**: 419–36.
- 18 Curley JP, Champagne FA. Influence of maternal care on the developing brain: mechanisms, temporal dynamics, and sensitive periods. *Front Neuroendocrinol* 2016; **40**: 52–66.

- 19 van IJzendoorn MH, Bakermans-Kranenburg MJ, Duschinsky R, et al. Institutionalisation and deinstitutionalisation of children 1: a systematic and integrative review of evidence regarding effects on development. *Lancet Psychiatry* 2020; **7**: 703–20.
- 20 Brett ZH, Humphreys KL, Fleming AS, Kraemer GW, Drury SS. Using cross-species comparisons and a neurobiological framework to understand early social deprivation effects on behavioral development. *Dev Psychopathol* 2015; **27**: 347–67.
- 21 Cameron JL, Eagleson KL, Fox NA, Hensch TK, Levitt P. Social origins of developmental risk for mental and physical illness. *J Neurosci* 2017; **37**: 10783–91.
- 22 Knudsen EI. Sensitive periods in the development of the brain and behavior. *J Cogn Neurosci* 2004; **16**: 1412–25.
- 23 Cicchetti D, Rogosch FA. Equifinality and multifinality in developmental psychopathology. *Dev Psychopathol* 1996; **8**: 597–600.
- 24 Page MJ, Moher D, Bossuyt PM, et al. PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. *BMJ* 2021; **372**: n160.
- 25 Cuthbert BN, Kozak MJ. Constructing constructs for psychopathology: the NIMH research domain criteria. *J Abnorm Psychol* 2013; **122**: 928–37.
- 26 Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan—a web and mobile app for systematic reviews. *Syst Rev* 2016; **5**: 210.
- 27 National Heart, Lung, and Blood Institute. Study quality assessment tools. 2021. <https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools> (accessed Dec 1, 2021).
- 28 Moran P, Eckenrode J. Protective personality characteristics among adolescent victims of maltreatment. *Child Abuse Negl* 1992; **16**: 743–54.
- 29 Lundqvist G, Hansson K, Svedin CG. The influence of childhood sexual abuse factors on women's health. *Nord J Psychiatry* 2004; **58**: 395–401.
- 30 Maercker A, Michael T, Fehm L, Becker ES, Margraf J. Age of traumatisation as a predictor of post-traumatic stress disorder or major depression in young women. *Br J Psychiatry* 2004; **184**: 482–7.
- 31 Schoedl AF, Costa MCP, Mari JJ, et al. The clinical correlates of reported childhood sexual abuse: an association between age at trauma onset and severity of depression and PTSD in adults. *J Child Sex Abuse* 2010; **19**: 156–70.
- 32 Gal G, Levav I, Gross R. Psychopathology among adults abused during childhood or adolescence: results from the Israel-based World Mental Health Survey. *J Nerv Ment Dis* 2011; **199**: 222–9.
- 33 Baker LM, Williams LM, Korgaonkar MS, Cohen RA, Heaps JM, Paul RH. Impact of early vs. late childhood early life stress on brain morphometrics. *Brain Imaging Behav* 2013; **7**: 196–203.
- 34 Alameda L, Golay P, Baumann PS, Ferrari C, Do KQ, Conus P. Age at the time of exposure to trauma modulates the psychopathological profile in patients with early psychosis. *J Clin Psychiatry* 2016; **77**: e612–18.
- 35 Alameda L, Golay P, Baumann PS, et al. Mild depressive symptoms mediate the impact of childhood trauma on long-term functional outcome in early psychosis patients. *Schizophr Bull* 2017; **43**: 1027–35.
- 36 Spak L, Spak F, Allebeck P. Sexual abuse and alcoholism in a female population. *Addiction* 1998; **93**: 1365–73.
- 37 Murphy SM, Kilpatrick DG, Amick-McMullan A, et al. Current psychological functioning of child sexual assault survivors: a community study. *J Interpers Violence* 1988; **3**: 55–79.
- 38 Thornberry TP, Ireland TO, Smith CA. The importance of timing: the varying impact of childhood and adolescent maltreatment on multiple problem outcomes. *Dev Psychopathol* 2001; **13**: 957–79.
- 39 Kaplow JB, Widom CS. Age of onset of child maltreatment predicts long-term mental health outcomes. *J Abnorm Psychol* 2007; **116**: 176–87.
- 40 Dunn EC, McLaughlin KA, Slopen N, Rosand J, Smoller JW. Developmental timing of child maltreatment and symptoms of depression and suicidal ideation in young adulthood: results from the National Longitudinal Study on Adolescent Health. *Depress Anxiety* 2013; **30**: 955–64.
- 41 Mueller-Pfeiffer C, Moergeli H, Schumacher S, et al. Characteristics of child maltreatment and their relation to dissociation, posttraumatic stress symptoms, and depression in adult psychiatric patients. *J Nerv Ment Dis* 2013; **201**: 471–77.
- 42 Harpur LJ, Polek E, van Harmelen AL. The role of timing of maltreatment and child intelligence in pathways to low symptoms of depression and anxiety in adolescence. *Child Abuse Negl* 2015; **47**: 24–37.
- 43 Dahl SK, Larsen JT, Petersen L, et al. Early adversity and risk for moderate to severe unipolar depressive disorder in adolescence and adulthood: a register-based study of 978,647 individuals. *J Affect Disord* 2017; **214**: 122–29.
- 44 Dunn EC, Nishimi K, Powers A, Bradley B. Is developmental timing of trauma exposure associated with depressive and post-traumatic stress disorder symptoms in adulthood? *J Psychiatr Res* 2017; **84**: 119–27.
- 45 Capretto JJ. Developmental timing of childhood physical and sexual maltreatment predicts adult depression and post-traumatic stress symptoms. *J Interpers Violence* 2017; **35**: 2558–82.
- 46 Seib C, McCarthy A, McGuire A, et al. Exposure to stress across the life course and its association with anxiety and depressive symptoms: results from the Australian Women's Wellness After Cancer Program (WWACP). *Maturitas* 2017; **105**: 107–12.
- 47 Atzl VM, Narayan AJ, Rivera LM, Lieberman AF. Adverse childhood experiences and prenatal mental health: type of ACEs and age of maltreatment onset. *J Fam Psychol* 2019; **33**: 304–14.
- 48 Merrick JS, Narayan AJ, Atzl VM, Harris WW, Lieberman AF. Type versus timing of adverse and benevolent childhood experiences for pregnant women's psychological and reproductive health. *Child Youth Serv Rev* 2020; **114**: 105056.
- 49 Carlin AS, Ward NG. Subtypes of psychiatric inpatient women who have been sexually abused. *J Nerv Ment Dis* 1992; **180**: 392–97.
- 50 Kumar G, Steer RA, Deblinger E. Problems in differentiating sexually from nonsexually abused adolescent psychiatric inpatients by self-reported anxiety, depression, internalization, and externalization. *Child Abuse Negl* 1996; **20**: 1079–86.
- 51 Yen S, Shea MT, Battle CL, et al. Traumatic exposure and posttraumatic stress disorder in borderline, schizotypal, avoidant, and obsessive-compulsive personality disorders: findings from the collaborative longitudinal personality disorders study. *J Nerv Ment Dis* 2002; **190**: 510–18.
- 52 Crane C, Duggan DS. Overgeneral autobiographical memory and age of onset of childhood sexual abuse in patients with recurrent suicidal behaviour. *Br J Clin Psychol* 2009; **48**: 93–100.
- 53 Khan A, McCormack HC, Bolger EA, et al. Childhood maltreatment, depression, and suicidal ideation: critical importance of parental and peer emotional abuse during developmental sensitive periods in males and females. *Front Psychiatry* 2015; **6**: 42.
- 54 Schalinski I, Teicher MH, Nischk D, Hinderer E, Müller O, Rockstroh B. Type and timing of adverse childhood experiences differentially affect severity of PTSD, dissociative and depressive symptoms in adult inpatients. *BMC Psychiatry* 2016; **16**: 295.
- 55 Gerke J, Koenig AM, Conrad D, et al. Childhood maltreatment as risk factor for lifetime depression: the role of different types of experiences and sensitive periods. *Ment Health Prev* 2018; **10**: 56–65.
- 56 Fujisawa TX, Shimada K, Takiguchi S, et al. Type and timing of childhood maltreatment and reduced visual cortex volume in children and adolescents with reactive attachment disorder. *NeuroImage Clin* 2018; **20**: 216–21.
- 57 Humphreys KL, King LS, Sacchet MD, et al. Evidence for a sensitive period in the effects of early life stress on hippocampal volume. *Dev Sci* 2019; **22**: e12775.
- 58 Luby JL, Tillman R, Barch DM. Association of timing of adverse childhood experiences and caregiver support with regionally specific brain development in adolescents. *JAMA Netw Open* 2019; **2**: e1911426.
- 59 Andersen SL, Tomada A, Vinco ES, Valente E, Polcari A, Teicher MH. Preliminary evidence for sensitive periods in the effect of childhood sexual abuse on regional brain development. *J Neuropsychiatry Clin Neurosci* 2008; **20**: 292–301.
- 60 Pechtel P, Lyons-Ruth K, Anderson CM, Teicher MH. Sensitive periods of amygdala development: the role of maltreatment in preadolescence. *NeuroImage Amst* 2014; **97**: 236–44.
- 61 Teicher MH, Anderson CM, Ohashi K, et al. Differential effects of childhood neglect and abuse during sensitive exposure periods on male and female hippocampus. *NeuroImage* 2018; **169**: 443–52.

- 62 Herzog JI, Thome J, Demirakca T, et al. Influence of severity of type and timing of retrospectively reported childhood maltreatment on female amygdala and hippocampal volume. *Sci Rep* 2020; **10**: 1903.
- 63 Enlow MB, Egeland B, Blood EA, Wright RO, Wright RJ. Interpersonal trauma exposure and cognitive development in children to age 8 years: a longitudinal study. *J Epidemiol Community Health* 2012; **66**: 1005–10.
- 64 Bell MF, Bayliss DM, Glauert R, Ohan JL. School readiness of maltreated children: associations of timing, type, and chronicity of maltreatment. *Child Abuse Negl* 2018; **76**: 426–39.
- 65 Jaffee SR, Maikovich-Fong AK. Effects of chronic maltreatment and maltreatment timing on children's behavior and cognitive abilities. *J Child Psychol Psychiatry* 2011; **52**: 184–94.
- 66 Dunn EC, Busso DS, Raffeld MR, et al. Does developmental timing of exposure to child maltreatment predict memory performance in adulthood? Results from a large, population-based sample. *Child Abuse Negl* 2016; **51**: 181–91.
- 67 Green MJ, Tzoumakis S, McIntyre B, et al. Childhood maltreatment and early developmental vulnerabilities at age 5 years. *Child Dev* 2018; **89**: 1599–612.
- 68 Holmes MR, Yoon S, Berg KA, Cage JL, Perzynski AT. Promoting the development of resilient academic functioning in maltreated children. *Child Abuse Negl* 2018; **75**: 92–103.
- 69 Hambrick EP, Brawner TW, Perry BD. Timing of early-life stress and the development of brain-related capacities. *Front Behav Neurosci* 2019; **13**: 183.
- 70 Rossen L, Tzoumakis S, Kariuki M, et al. Timing of the first report and highest level of child protection response in association with early developmental vulnerabilities in an Australian population cohort. *Child Abuse Negl* 2019; **93**: 1–12.
- 71 Mackiewicz Seghete KL, DePrince AP, Banich MT. Association between initial age of exposure to childhood abuse and cognitive control: preliminary evidence. *J Trauma Stress* 2018; **31**: 437–47.
- 72 Schalinski I, Teicher MH, Carolus AM, Rockstroh B. Defining the impact of childhood adversities on cognitive deficits in psychosis: an exploratory analysis. *Schizophr Res* 2018; **192**: 351–56.
- 73 Jun HJ, Rich-Edwards JW, Boynton-Jarrett R, Austin SB, Frazier AL, Wright RJ. Child abuse and smoking among young women: the importance of severity, accumulation, and timing. *J Adolesc Health* 2008; **43**: 55–63.
- 74 Fisher HL, Jones PB, Fearon P, et al. The varying impact of type, timing and frequency of exposure to childhood adversity on its association with adult psychotic disorder. *Psychol Med* 2010; **40**: 1967–78.
- 75 McCutcheon VV, Sartor CE, Pommer NE, et al. Age at trauma exposure and PTSD risk in young adult women. *J Trauma Stress* 2010; **23**: 811–14.
- 76 Mossige S, Huang L, Straiton M, Roen K. Suicidal ideation and self-harm among youths in Norway: associations with verbal, physical and sexual abuse. *Child Fam Soc Work* 2016; **21**: 166–75.
- 77 Yoon S, Pei F, Wang X, et al. Vulnerability or resilience to early substance use among adolescents at risk: the roles of maltreatment and father involvement. *Child Abuse Negl* 2018; **86**: 206–16.
- 78 Manly JT, Kim JE, Rogosch FA, Cicchetti D. Dimensions of child maltreatment and children's adjustment: contributions of developmental timing and subtype. *Dev Psychopathol* 2001; **13**: 759–82.
- 79 Kotch JB, Lewis T, Hussey JM, et al. Importance of early neglect for childhood aggression. *Pediatrics* 2008; **121**: 725–31.
- 80 Yoon S, Dillard R, Kobulsky J, Nemeth J, Shi Y, Schoppe-Sullivan S. The type and timing of child maltreatment as predictors of adolescent cigarette smoking trajectories. *Subst Use Misuse* 2020; **55**: 937–46.
- 81 Chu JA, Frey LM, Ganzel BL, Matthews JA. Memories of childhood abuse: dissociation, amnesia, and corroboration. *Am J Psychiatry* 1999; **156**: 749–55.
- 82 Schalinski I, Breinlinger S, Hirt V, Teicher MH, Odenwald M, Rockstroh B. Environmental adversities and psychotic symptoms: the impact of timing of trauma, abuse, and neglect. *Schizophr Res* 2019; **205**: 4–9.
- 83 Teicher MH, Parigger A. The 'Maltreatment and Abuse Chronology of Exposure' (MACE) scale for the retrospective assessment of abuse and neglect during development. *PLoS One* 2015; **10**: e0117423.
- 84 Kaczurkin AN, Raznahan A, Satterthwaite TD. Sex differences in the developing brain: insights from multimodal neuroimaging. *Neuropsychopharmacology* 2019; **44**: 71–85.
- 85 Dunn EC, Gilman SE, Willett JB, Slopen NB, Molnar BE. The impact of exposure to interpersonal violence on gender differences in adolescent-onset major depression: results from the National Comorbidity Survey Replication (NCS-R). *Depress Anxiety* 2012; **29**: 392–99.
- 86 Zhu Y, Wang MJ, Crawford KM, et al. Sensitive period-regulating genetic pathways and exposure to adversity shape risk for depression. *Neuropsychopharmacology* 2022; **47**: 497–506.
- 87 Batista G, Hensch TK. Critical period regulation by thyroid hormones: potential mechanisms and sex-specific aspects. *Front Mol Neurosci* 2019; **12**: 77.
- 88 Premachandran H, Zhao M, Arruda-Carvalho M. Sex differences in the development of the rodent corticolimbic system. *Front Neurosci* 2020; **14**: 583477.
- 89 Castrén E, Antila H. Neuronal plasticity and neurotrophic factors in drug responses. *Mol Psychiatry* 2017; **22**: 1085–95.
- 90 Kumsta R, Stevens S, Brookes K, et al. 5HTT genotype moderates the influence of early institutional deprivation on emotional problems in adolescence: evidence from the English and Romanian Adoptee (ERA) study. *J Child Psychol Psychiatry* 2010; **51**: 755–62.
- 91 Zeanah CH, Egger HL, Smyke AT, et al. Institutional rearing and psychiatric disorders in Romanian preschool children. *Am J Psychiatry* 2009; **166**: 777–85.
- 92 Humphreys KL, Gleason MM, Drury SS, et al. Effects of institutional rearing and foster care on psychopathology at age 12 years in Romania: follow-up of an open, randomised controlled trial. *Lancet Psychiatry* 2015; **2**: 625–34.
- 93 Sheridan MA, McLaughlin KA. Dimensions of early experience and neural development: deprivation and threat. *Trends Cogn Sci* 2014; **18**: 580–85.
- 94 Sheridan MA, Peverill M, Finn AS, McLaughlin KA. Dimensions of childhood adversity have distinct associations with neural systems underlying executive functioning. *Dev Psychopathol* 2017; **29**: 1777–94.
- 95 Pechtel P, Lyons-Ruth K, Anderson CM, Teicher MH. Sensitive periods of amygdala development: the role of maltreatment in preadolescence. *Neuroimage* 2014; **97**: 236–44.
- 96 Kaplow JB, Widom CS. Age of onset of child maltreatment predicts long-term mental health outcomes. *J Abnorm Psychol* 2007; **116**: 176–87.
- 97 Dunn EC, McLaughlin KA, Slopen N, Rosand J, Smoller JW. Developmental timing of child maltreatment and symptoms of depression and suicidal ideation in young adulthood: results from the National Longitudinal Study of Adolescent Health. *Depress Anxiety* 2013; **30**: 955–64.
- 98 Weissman DG, Bitran D, Miller AB, Schaefer JD, Sheridan MA, McLaughlin KA. Difficulties with emotion regulation as a transdiagnostic mechanism linking child maltreatment with the emergence of psychopathology. *Dev Psychopathol* 2019; **31**: 899–915.
- 99 Vachon DD, Krueger RF, Rogosch FA, Cicchetti D. Assessment of the harmful psychiatric and behavioral effects of different forms of child maltreatment. *JAMA Psychiatry* 2015; **72**: 1135–42.
- 100 Reuben A, Moffitt TE, Caspi A, et al. Lest we forget: comparing retrospective and prospective assessments of adverse childhood experiences in the prediction of adult health. *J Child Psychol Psychiatry* 2016; **57**: 1103–12.
- 101 Newbury JB, Arseneault L, Moffitt TE, et al. Measuring childhood maltreatment to predict early-adult psychopathology: comparison of prospective informant-reports and retrospective self-reports. *J Psychiatr Res* 2018; **96**: 57–64.
- 102 Baldwin JR, Reuben A, Newbury JB, Danese A. Agreement between prospective and retrospective measures of childhood maltreatment: a systematic review and meta-analysis. *JAMA Psychiatry* 2019; **76**: 584–93.
- 103 Simon GE, VonKorff M. Recall of psychiatric history in cross-sectional surveys: implications for epidemiologic research. *Epidemiol Rev* 1995; **17**: 221–27.
- 104 Danese A. Annual research review: rethinking childhood trauma—new research directions for measurement, study design and analytical strategies. *J Child Psychol Psychiatry* 2020; **61**: 236–50.

- 105 Davis KA, Mountain RV, Pickett OR, Den Besten PK, Bidlack FB, Dunn EC. Teeth as potential new tools to measure early-life adversity and subsequent mental health risk: an interdisciplinary review and conceptual model. *Biol Psychiatry* 2020; **87**: 502–13.
- 106 Mountain RV, Zhu Y, Pickett OR, et al. Association of maternal stress and social support during pregnancy with growth marks in children's primary tooth enamel. *JAMA Netw Open* 2021; **4**: e2129129.
- 107 Dunn EC, Soare TW, Raffeld MR, et al. What life course theoretical models best explain the relationship between exposure to childhood adversity and psychopathology symptoms: recency, accumulation, or sensitive periods? *Psychol Med* 2018; **48**: 2562–72.
- 108 Smith BJ, Smith ADAC, Dunn EC. Statistical modeling of sensitive period effects using the Structured Life Course Modeling Approach (SLCMA). *Curr Top Behav Neurosci* 2022; **53**: 215–34.
- 109 Zhu J, Lowen SB, Anderson CM, Ohashi K, Khan A, Teicher MH. Association of prepubertal and postpubertal exposure to childhood maltreatment with adult amygdala function. *JAMA Psychiatry* 2019; **76**: 843–53.
- 110 Simonsohn U, Simmons JP, Nelson LD. Specification curve analysis. *Nat Hum Behav* 2020; **4**: 1208–14.
- 111 Ben-Shlomo Y, Kuh D. A life course approach to chronic disease epidemiology: conceptual models, empirical challenges and interdisciplinary perspectives. *Int J Epidemiol* 2002; **31**: 285–93.
- 112 Cicchetti D. Neural plasticity, sensitive periods, and psychopathology. *Dev Psychopathol* 2015; **27**: 319–20.

Copyright © 2022 Elsevier Ltd. All rights reserved.